

ORIGINAL ARTICLE

Detection of Extended-Spectrum β -Lactamases among *Acinetobacter Baumannii* Isolated from Hospitals of Qazvin, Iran**Mina Zarabadi-Pour¹, Amir Peymani^{2*}, Narges Habibollah-Pourzereshki², Mohammad Reza Sarookhani³, Ali Akbar Karami⁴, Amir Javadi²**

OPEN ACCESS

Citation: Mina Zarabadi-Pour, Amir Peymani, Narges Habibollah-Pourzereshki, Mohammad Reza Sarookhani, Ali Akbar Karami, Amir Javadi. Detection of Extended-Spectrum β -Lactamases among *Acinetobacter Baumannii* Isolated from Hospitals of Qazvin, Iran. *Ethiop J Health Sci*. 2021;31 (2):229. doi:<http://dx.doi.org/10.4314/ejhs.v31i2.4>

Received: October 9, 2020

Accepted: October 22, 2020

Published: March 1, 2021

Copyright: © 2021 Mina Z., et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This study was funded by the Research Deputy of Qazvin University of Medical Sciences (Grant No. 10016).

Competing Interests: The authors declare that this manuscript was approved by all authors in its form and that no competing interest exists.

Affiliation and Correspondence:

¹Student Research Committee, Qazvin University of Medical Sciences, Qazvin, Iran

²Medical Microbiology Research Center, Qazvin University of Medical Sciences, Qazvin, Iran

³Cellular and Molecular Research Center, Qazvin University of Medical Sciences, Qazvin, Iran

⁴Department of Urology, School of Medicine, Qazvin University of Medical Sciences, Qazvin, Iran

*Email: a.peymani@gmail.com

ABSTRACT

BACKGROUND: *Acinetobacter baumannii* is a major contributor to nosocomial infections. Extended-spectrum β -lactamase (ESBL)-producing *A. baumannii* is spreading worldwide. We aimed to determine the frequency of ESBL-encoding genes in clinical isolates of *A. baumannii* and to access their clonal relationship by repetitive extragenic palindromic-PCR (rep-PCR).

METHODS: In this descriptive cross-sectional study, 203 isolates of *A. baumannii* were collected from Qazvin hospitals. The Identification of isolates was performed by standard laboratory methods. To verify ESBL production, all isolates were screened by disk agar diffusion and confirmed by the combined disk method. Subsequently, ESBL-encoding genes were detected by PCR and sequencing. Possible clonal association of ESBL-producing isolates was evaluated using rep-PCR.

RESULTS: Two hundred (98.5%) isolates showed reduced susceptibility to one of the antibiotics used in the ESBL screening test, of which 127 isolates (62.6%) produced ESBL. PCR results showed *bla*_{OXA-1} (20.5%) was the most prevalent gene followed by *bla*_{TEM-1} (20%), *bla*_{GES-1} (15.7%), *bla*_{CTX-M-15} (7.9%), and *bla*_{PER-1} (1.6%). Rep-PCR results revealed that ESBL-producing isolates belonged to clones A (85%), B (13.4%), and C (1.6%).

CONCLUSION: Our study showed the significant presence of *bla*_{OXA-1}, *bla*_{TEM-1}, *bla*_{GES-1}, *bla*_{CTX-M-15}, and *bla*_{PER-1} genes in ESBL-producing *A. baumannii* isolates in the studied hospitals. This is the first report on the emergence of *bla*_{OXA-1} gene in these isolates in Iran. The use of comprehensive antimicrobial treatment guidelines based on laboratory data and appropriate infection control interventions are essential.

KEYWORDS: *Acinetobacter baumannii*, ESBL, repetitive extragenic palindromic-PCR